

# Cilatil

**QUICK GUIDE<sup>1</sup>**

10 mg & 20 mg

## WHEN TO TAKE IT

At least **30 - 60 minutes** prior to anticipated sexual activity

---

## WHEN WILL IT START WORKING

Varies from person to person

From **30 minutes**

---

## HOW LONG WILL IT WORK FOR

Works only when sexually aroused

Up to **36 hours** when you need it

## FOOD AND ALCOHOL

Drinking alcohol may affect your ability to get an erection, so avoid excessive drinking

**Yes**, can be taken with a meal

---

## FREQUENCY OF USE

Not for continuous daily use.  
Talk to your GP for alternatives if taking twice weekly or more

---

## SIDE EFFECTS\*

Headache, indigestion, back pain, sore throat, muscle pain, decrease or loss of hearing



**CILATIL<sup>®</sup>**  
tadalafil

Scan QR code  
for more information



[cilatil.co.nz](http://cilatil.co.nz)

## CILATIL REWARDS

Enjoy every

5<sup>th</sup> PACK 

Terms and conditions to redeem 5<sup>th</sup> pack apply

Scan QR code for more information



[cilatil.co.nz/en/cilatil-rewards](https://cilatil.co.nz/en/cilatil-rewards)

\*These are some of the more common or/and very serious side effects. For a comprehensive list consult your healthcare professional or refer to the Consumer Medicine Information for more details.

**References:** 1. Cilatil<sup>®</sup> Consumer Medicine Information (4 May 2023). Available at <https://www.medsafe.govt.nz>. Accessed March 2024. **Cilatil<sup>®</sup>** (tadalafil) 5 mg, 10 mg & 20 mg tablets. Prescription Medicine. For the treatment of adult men with erectile dysfunction (ED), moderate to severe lower urinary tract symptoms (LUTS) associated with benign prostatic hyperplasia (BPH), with coexisting ED and LUTS associated with BPH. Do not take if you: are allergic to tadalafil or to any excipients; taking nitrates or guanylate cyclase stimulators, have a cardiac disease for which sexual activity is inadvisable, have uncontrolled arrhythmias or blood pressure, previous eye condition called NAION. Medicines have benefits and risks. Possible side effects headache, indigestion, back pain, muscle pain. Serious side effects may include: prolonged or painful erection, decrease or loss of vision. Take strictly as directed. If symptoms persist or you have side effects, see your healthcare professional. Ask your doctor if Cilatil<sup>®</sup> is right for you. For more information, view the Consumer Medicine Information at [www.medsafe.govt.nz](https://www.medsafe.govt.nz). Cilatil<sup>®</sup> is an unfunded medicine - doctor's fees and pharmacy charges will apply. Cilatil<sup>®</sup> is a Viatrix company trade mark, Viatrix Limited, Auckland. Copyright© 2024 Viatrix Inc. All rights reserved. TAD-2024-0015. TAPS NP20642.

QUICK GUIDE<sup>1</sup>



**CILATIL<sup>®</sup>**  
tadalafil

Ready when you are, again and again and again...  
Hope for couples with erection problems.



**VIATRIS**